



OPEN ACCESS

EDITED BY

Silvia Pogliaghi,
University of Verona, Italy

REVIEWED BY

Enrico Tam,
University of Verona, Italy
Giovanni Vinetti,
Eurac Research, Italy

*CORRESPONDENCE

Andrea Nicolò,
✉ andrea.nicolo@uniroma4.it

RECEIVED 21 May 2023

ACCEPTED 20 July 2023

PUBLISHED 01 August 2023

CITATION

Nicolò A, Girardi M, Bazzucchi I,
Sacchetti M and Felici F (2023),
Ventilation and perceived exertion are
sensitive to changes in exercise
tolerance: arm+leg cycling vs.
leg cycling.
Front. Physiol. 14:1226421.
doi: 10.3389/fphys.2023.1226421

COPYRIGHT

© 2023 Nicolò, Girardi, Bazzucchi,
Sacchetti and Felici. This is an open-
access article distributed under the terms
of the [Creative Commons Attribution
License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or
reproduction in other forums is
permitted, provided the original author(s)
and the copyright owner(s) are credited
and that the original publication in this
journal is cited, in accordance with
accepted academic practice. No use,
distribution or reproduction is permitted
which does not comply with these terms.

Ventilation and perceived exertion are sensitive to changes in exercise tolerance: arm+leg cycling vs. leg cycling

Andrea Nicolò ^{1*}, Michele Girardi ^{1,2}, Ilenia Bazzucchi ¹,
Massimo Sacchetti ¹ and Francesco Felici ¹

¹Laboratory of Exercise Physiology, Department of Movement, Human and Health Sciences, University of Rome "Foro Italico", Rome, Italy, ²Division of Respiratory and Critical Care Physiology and Medicine, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Institute of Respiratory Medicine and Exercise Physiology, Torrance, CA, United States

Purpose: Growing evidence suggests that respiratory frequency (f_R) is a marker of physical effort and a variable sensitive to changes in exercise tolerance. The comparison between arm+leg cycling (Arm+leg) and leg cycling (Leg) has the potential to further test this notion because a greater exercise tolerance is expected in the Arm+leg modality. We systematically compared Arm+leg vs. Leg using different performance tests.

Methods: Twelve males underwent six performance tests in separate, randomized visits. Three tests were performed in each of the two exercise modalities, i.e. an incremental test and two time-to-exhaustion (TTE) tests performed at 90% or 75% of the peak power output reached in the Leg incremental test (PPO_{Leg}). Exercise tolerance, perceived exertion, and cardiorespiratory variables were recorded during all the tests.

Results: A greater exercise tolerance ($p < 0.001$) was found for Arm+leg in the incremental test (337 ± 32 W vs. 292 ± 28 W), in the TTE test at 90% of PPO_{Leg} (638 ± 154 s vs. 307 ± 67 s), and in the TTE test at 75% of PPO_{Leg} ($1,675 \pm 525$ s vs. 880 ± 363 s). Unlike $\dot{V}O_2$ and heart rate, both f_R and minute ventilation were lower ($p < 0.003$) at isotime in all the Arm+leg tests vs. Leg tests. Furthermore, a lower perceived exertion was observed in the Arm+leg tests, especially during the TTE tests ($p < 0.001$).

Conclusion: Minute ventilation, f_R and perceived exertion are sensitive to the improvements in exercise tolerance observed when comparing Arm+leg vs. Leg, unlike $\dot{V}O_2$ and heart rate.

Abbreviations: Arm+leg, arm+leg cycling; Arm+leg_{INC}, arm+leg cycling incremental test; Arm+leg_{TTE75}, Arm+leg TTE test at 75% of PPO_{Leg} ; Arm+leg_{TTE90}, Arm+leg TTE test at 90% of PPO_{Leg} ; f_R , respiratory frequency; GET, gas exchange threshold; HR, heart rate; Leg, leg cycling; Leg_{INC}, leg cycling incremental test; Leg_{TTE75}, Leg TTE test at 75% of PPO_{Leg} ; Leg_{TTE90}, Leg TTE test at 90% of PPO_{Leg} ; P_{ETCO_2} , end-tidal partial pressure of carbon dioxide; PPO, Peak power output; PPO_{Leg} , peak power output reached in the Leg_{INC}; RCP, respiratory compensation point; RPE, ratings of perceived exertion; TTE, time-to-exhaustion; $\dot{V}CO_2$, carbon dioxide output; \dot{V}_E , minute ventilation; $\dot{V}O_2$, oxygen uptake; $\dot{V}O_{2peak}$, peak value of oxygen uptake; V_T , tidal volume.

KEYWORDS

endurance performance, breathing control, respiratory frequency, incremental test, time to exhaustion, fatigue, breathing pattern, oxygen uptake

1 Introduction

The comparison between different exercise modalities has the potential to improve our understanding of the physiology of endurance performance. Classical studies have compared leg cycling (Leg) with arm+leg cycling (Arm+leg) to unravel the mechanisms limiting maximal oxygen uptake (Astrand and Saltin, 1961; Secher et al., 1974; Bergh et al., 1976). Findings from these studies have contributed to outlining the important role of cardiocirculatory factors in setting the upper limit for maximal aerobic power. Indeed, the peak value of oxygen uptake ($\dot{V}O_{2\text{peak}}$) is not always proportional to the differences in the amount of muscle mass involved in various exercise modalities, as the addition of arm work to leg work generally does not increase $\dot{V}O_{2\text{peak}}$ more than about 5%–10% (Astrand and Saltin, 1961; Gleser et al., 1974; Secher et al., 1974; Secher and Volianitis, 2006). In fact, the comparison between arm+leg cycling and leg cycling is suitable for gaining insight into other physiological responses that have received less attention so far, including the variables associated with physical effort and changes in exercise tolerance. Exercise tolerance is here defined as the tolerated duration during a time-to-exhaustion (TTE) test performed at a constant work rate or the peak power output (PPO) achieved during an incremental test (Van De Walle and Vukovich, 2018). Some findings have shown that arm+leg cycling results in a greater exercise tolerance compared to leg cycling alone (arms hanging on the participant's side) (Gleser et al., 1974; Secher et al., 1974; Nagle et al., 1984), but it is unclear if this difference is still evident when arm+leg cycling is compared to conventional leg cycling (Secher et al., 1974; Bergh et al., 1976; Nagle et al., 1984). When matched for the same absolute total power output, the greater exercise tolerance that might be expected for arm+leg cycling makes the comparison with (conventional) leg cycling valuable for testing the proposition that improvements in exercise tolerance are accompanied by consistent changes in the responses of respiratory frequency (f_R) and perceived exertion (Nicolò and Sacchetti, 2023).

Growing evidence suggests that f_R is a valid marker of physical effort (Nicolò et al., 2014; Nicolò et al., 2016; Nicolò et al., 2017a; Nicolò et al., 2017b; Nicolò et al., 2019; Nicolò and Sacchetti, 2023) and that its time course reflects changes in exercise tolerance (Nicolò and Sacchetti, 2023). In a variety of conditions where exercise tolerance is reduced (experimentally) or lowered (in a cross-sectional comparison), the rate of increase in f_R is higher, both during incremental and TTE tests. On the other hand, the rate of increase in f_R is lower when assessing exercise strategies, experimental interventions or other conditions leading to an improvement in exercise tolerance (Nicolò and Sacchetti, 2023). The sensitivity of f_R to changes in exercise tolerance and the close association between f_R and perceived exertion are among the factors suggesting that f_R is to a large extent modulated by central command (the activity of motor and premotor brain areas relating to voluntary locomotor muscle contraction) during high-intensity exercise (Nicolò and Sacchetti, 2023). This explains why f_R can be considered a marker of physical effort, which is defined as the

degree of motor effort (i.e. the magnitude of central command) (Nicolò et al., 2017b). However, measuring the magnitude of central command during “real” exercise conditions is particularly challenging. Hence, it is important to use different approaches (including the comparison of different exercise modalities) to provide indirect evidence on the contribution of central command to f_R modulation (Nicolò and Sacchetti, 2023).

The comparison between arm+leg cycling and leg cycling may either challenge or reinforce the notion that the increase in f_R during high-intensity exercise reflects changes in exercise tolerance and is influenced by central command. f_R is also modulated by muscle afferent feedback from groups III and IV (hereinafter muscle afferent feedback) (Dempsey et al., 2014; Girardi et al., 2021; Nicolò and Sacchetti, 2023), and this drive to breathe may have a greater relative contribution to f_R modulation when arm muscles assist leg muscles during arm+leg cycling, in view of a potentially larger amount of muscle mass concomitantly involved in exercise (Dempsey et al., 2014; Nicolò and Sacchetti, 2023). Indeed, even the passive movement of the legs leads to a substantial increase in f_R that is at least partially mediated by muscle afferent feedback (Girardi et al., 2021), and this drive to breathe may increase further when adding the movement of the upper limbs. The contribution of muscle afferent feedback to f_R may partially confound the association between f_R and perceived exertion because the latter is supposed to be largely independent of muscle afferent feedback (Marcora, 2009; Bergevin et al., 2023). While arm+leg cycling may also result in a higher $\dot{V}O_2$ compared to leg cycling at the same submaximal power output (Hoffman et al., 1996), f_R largely dissociates from metabolic rate and is not substantially modulated by metabolic inputs, unlike tidal volume (V_T) (Nicolò et al., 2017a; 2018; Nicolò and Sacchetti, 2019; Nicolò et al., 2020a; Nicolò and Sacchetti, 2023). On the other hand, some findings seem to support the association between f_R and perceived exertion during both arm+leg cycling and leg cycling. Robertson et al. (1986) found similar responses of f_R —but not V_T —and the ratings of perceived exertion (RPE) during arm cycling, leg cycling and arm+leg cycling for intensities ranging from 20% to 80% of $\dot{V}O_{2\text{peak}}$. However, the authors neither reported the responses of f_R and RPE when exhaustion was approaching nor described if the two variables were sensitive to between-modality changes in exercise tolerance. Further studies are required to address this issue.

The purpose of the present study was to systematically assess whether exercise tolerance improves with arm+leg cycling vs. (conventional) leg cycling and whether f_R and perceived exertion are sensitive to the expected differences in exercise tolerance. To increase the robustness of our evaluation, we compared arm+leg cycling vs. leg cycling using two exercise paradigms (i.e. incremental test and TTE test) and three comparisons, as the TTE test was performed at two different intensities. We tested the hypotheses that i) arm+leg cycling improves exercise tolerance compared to leg cycling irrespective of the exercise paradigm; and ii) f_R is a good marker of physical effort sensitive to between-modality changes in exercise tolerance, unlike other physiological variables such as $\dot{V}O_2$ and heart rate (HR).




FIGURE 1

Three-dimensional (A) and lateral (B) views of the multimodal ergometer composed of a grinding ergometer and a cycling ergometer. The multimodal ergometer was custom-made by ORF s.r.l Magnetic Days® (Arezzo, Italy). Note that each of the two ergometers is electromagnetically braked and is equipped with a professional torque transducer.

2 Materials and methods

2.1 Participants

Twelve recreationally trained males (mean \pm SD: age 26 ± 4 years; stature 1.79 ± 0.08 m and body mass: 81 ± 10 kg) volunteered to participate in this study. The volunteers recruited participated in one or more sporting activities requiring the use of both arms and legs (e.g. rugby, extreme conditioning program training, and triathlon), as the benefits of arm+leg cycling vs. leg cycling may be more pronounced for individuals exercising with both upper and lower limbs (Secher et al., 1974). The study was approved by the Institutional Review Board of the University of Rome “Foro Italico” in compliance with the *Declaration of Helsinki* (CAR 07/2019). Written informed consent was obtained from all participants. They were asked to refrain from vigorous exercise and the consumption of alcohol and caffeine in the 24 h preceding each laboratory visit.

2.2 Experimental overview

Participants reported to the laboratory on 7 different occasions over a 4-week period, with visits separated by at least 48 h. On the first visit, participants were familiarised with the experimental procedures and tests. On the subsequent visits, participants performed three performance tests to exhaustion in each of the two exercise modalities, i.e. arm+leg cycling and leg cycling. The performance tests consisted of a step incremental test and two TTE tests performed at different intensities. Specifically, the incremental tests (Arm+leg_{INC} and Leg_{INC}) were performed on visits 2 and 3, in random order. The PPO of the Leg_{INC} test (PPO_{Leg}) was used to set the power output of the TTE tests (i.e. 90% and 75% of PPO_{Leg}), which were performed on visits 4–7. The order of Arm+leg and Leg

tests was always randomized, as well as the order of the TTE tests at 90% and 75% of PPO_{Leg}. All the tests were performed on a multimodal ergometer custom-made by ORF s.r.l Magnetic Days® (Arezzo, Italy) and specifically developed for performing this study. Exhaustion was defined as the decrease in pedaling cadence below 60 rpm, either with the legs or arms. All testing was completed in a laboratory with a room temperature of $23^{\circ}\text{C} \pm 1^{\circ}\text{C}$. A cooling fan was used during all the tests, and mechanical, physiological, and perceptual variables were recorded as detailed below.

2.3 Multimodal ergometer

The multimodal ergometer was made up of a grinding ergometer and a cycling ergometer, and their combined use allowed the participants to perform arm+leg cycling (see Figure 1). Both the grinding and cycling ergometers were electromagnetically braked, equipped with professional torque transducers (model RT2A) certified by AEP transducers (Cognento, Modena, Italy), and calibrated according to the manufacturer’s instructions. The expanded uncertainty was lower than 0.114% for both torque transducers used. The arm cranks of the grinding ergometer were provided by Harken Italy SPA (Limido Comasco, Como). The chainring of the grinding ergometer was not mechanically connected with that of the cycling ergometer to allow for the separate measurement of the power output provided by the two ergometers. Hence, one of the advantages of this multimodal ergometer is the opportunity to set and register the contribution of arms and legs to the total power output. In all the Arm+leg tests, the relative contribution of the arms was initially set at 20% of the total power output based on the findings reported by Bergh et al. (1976). Thereafter, participants were allowed to request changes in the relative contribution of the arms throughout each test according to preference, and they were familiarised with this procedure on the

first visit. The option of individualizing the relative contribution of the arms is supported by previous studies reporting inter-individual variability in preference (Hill et al., 2018) and exercise tolerance (Bergh et al., 1976) for different power output distributions between arms and legs. Participants were free to choose their preferred pedaling cadence in all the tests. In both Arm+leg and Leg modalities, the ergometer settings were set up on the first visit according to participants' anthropometric characteristics and comfort, and were reproduced in the subsequent visits. The Arm+leg tests were performed on the multimodal ergometer, while the Leg tests were performed on the cycling ergometer (the participants were allowed to use the handlebars).

2.4 Step incremental tests

Before the incremental tests, a 5-min warm-up was performed to allow the participants to check the ergometer settings. For both Arm+leg_{INC} and Leg_{INC}, the first stage of the incremental test consisted of 3 min at 150 W, and the power output was subsequently increased by 20 W every min. The power output of the first stage was chosen to ensure that at least 20% of the total power output could be sustained by the arms in the Arm+leg modality, and the minimum power output for the arm ergometer was about 25 W. Hence, it was not possible to select a power output lower than 150 W, and this limited the opportunity to rigorously determine the gas exchange threshold (GET) and the respiratory compensation point (RCP). The PPO reached in the incremental test was computed as the work rate of the last completed stage plus the fraction of time spent in the last uncompleted stage multiplied by the work-rate increment (i.e. 20 W). The Borg's 6–20 scale (Borg, 1998) was used to collect ratings of perceived exertion (RPE) data every min. Participants were familiarised with the use of the RPE scale on the first visit and were asked to verbally provide an RPE value, as required by the Arm+leg modality. Breathing artifacts caused by speaking were then removed by data filtering, as described below. Participants did not receive any performance feedback or encouragement during any of the incremental tests performed.

2.5 Time to exhaustion tests

After a 10-min self-paced warm-up, participants performed a TTE test in each of the two exercise modalities (Arm+leg and Leg) and intensities (90% and 75% of PPO_{Leg}) in separate visits. Hereinafter, the TTE tests are abbreviated as Arm+leg_{TTE90} (Arm+leg test at 90% of PPO_{Leg}), Leg_{TTE90} (Leg test at 90% of PPO_{Leg}), Arm+leg_{TTE75} (Arm+leg test at 75% of PPO_{Leg}) and Leg_{TTE75} (Leg test at 75% of PPO_{Leg}). Perceived exertion data were collected every min, while physiological and mechanical variables were measured continuously. Participants did not receive any performance feedback or encouragement during any of the TTE tests performed in this study.

2.6 Cardiorespiratory measures

f_R , V_T , minute ventilation (\dot{V}_E), $\dot{V}O_2$, carbon dioxide output ($\dot{V}CO_2$), end-tidal partial pressure of carbon dioxide (P_{ETCO_2}) and

HR were measured breath-by-breath during all the tests using a metabolic cart (Quark CPET, Cosmed, Rome, Italy). The metabolic cart was calibrated following the manufacturer's instructions.

2.7 Data analysis

Data were analyzed with MATLAB (R2016a, The Mathworks, Natick, MA, United States). The comparison of the physiological and perceptual responses between arm+leg cycling and leg cycling was performed in all the tests using the "individual isotime" analysis described by Nicolò et al. (2019). This analysis allows for between-condition comparisons while avoiding the data loss that occurs when the variability in TTE is not addressed on an individual basis (Nicolò et al., 2019). Briefly, breath-by-breath data of f_R , V_T , \dot{V}_E , $\dot{V}O_2$, $\dot{V}CO_2$, P_{ETCO_2} and HR were filtered for errant breaths by deleting values greater than 3 standard deviations from the local mean (Lamarra et al., 1987). Subsequently, breath-by-breath data were linearly interpolated and extrapolated every second. Data were then smoothed by a moving average of 60 s. Likewise, RPE data collected every min were linearly interpolated and extrapolated every second. Thereafter, for each individual, the shortest test of each Arm+leg vs. Leg comparison (Arm+leg_{INC} vs. Leg_{INC}, Arm+leg_{TTE90} vs. Leg_{TTE90} and Arm+leg_{TTE75} vs. Leg_{TTE75} were compared separately) was segmented into ten timepoints, and the same segmentation was used for the longest test of the same participant. This procedure was performed for all the participants as further detailed by Nicolò et al. (2019).

When reporting the relationship between different variables (i.e. RPE vs. f_R , RPE vs. \dot{V}_E , RPE vs. HR, \dot{V}_E vs. V_T , \dot{V}_E vs. $\dot{V}CO_2$, V_T vs. $\dot{V}CO_2$, and f_R vs. $\dot{V}CO_2$), another analysis called "relative isotime" was used as previously suggested (Nicolò et al., 2019). This analysis segments each test into ten timepoints based on the TTE of the test analyzed, and thus results in no data loss for any of the tests.

2.8 Statistical analysis

Statistical analyses were conducted using IBM SPSS Statistics 20 (SPSS Inc., Chicago, IL, United States). Data were checked for normality prior to analysis. A paired Student's *t*-test was used to compare the performance values of Arm+leg_{INC} vs. Leg_{INC}, Arm+leg_{TTE90} vs. Leg_{TTE90} and Arm+leg_{TTE75} vs. Leg_{TTE75} separately. The Cohen's *d* effect size for paired *t*-test was then calculated and considered small, moderate or large for values ≥ 0.2 , ≥ 0.5 and ≥ 0.8 respectively. A paired Student's *t*-test was also used to compare the end-test values of physiological variables between Arm+leg and Leg tests. A two-way repeated-measures ANOVA (condition \times time) was used to compare physiological and perceptual responses (processed with the "individual isotime" method) of Arm+leg vs. Leg for each of the three performance tests separately. When the sphericity assumption was violated, the Greenhouse–Geisser adjustment was performed. Partial eta squared (η_p^2) effect sizes were calculated for the main effect of condition, the main effect of time, and the interaction; η_p^2 values ≥ 0.01 , ≥ 0.059 and ≥ 0.138 indicate small, medium and large effects respectively (Cohen, 1988). When a significant interaction was found, pairwise comparisons were performed at each time point




FIGURE 2

Average performance differences between Arm+leg tests (blue bar graphs) and Leg tests (grey bar graphs) for the incremental test (A), the TTE test at 90% of PPO_{Leg} (B), and the TTE test at 75% of PPO_{Leg} (C). The open circles and the filled circles represent individual data during Arm+leg and Leg tests respectively. * $p < 0.05$ vs. Arm+leg.

using a one-way repeated measures ANOVA to identify differences between Arm+leg and Leg tests. The HR data of the incremental tests and the TTE tests at 75% of PPO_{Leg} were not normally distributed and were analyzed using Friedman's two-way ANOVA. When statistical significance was found, this test was followed up by a Wilcoxon Signed-Rank test to identify where differences between Arm+leg and Leg tests occurred.

After processing data with the "relative isotime" method, the correlations between RPE and f_R , RPE and \dot{V}_E , and RPE and HR were analyzed using a previously described method that adjusts for repeated observations within participants (Bland and Altman, 1995). A correlation coefficient (r) and a p -value were obtained by considering all the performance tests together. A p -value < 0.05 was considered statistically significant in all analyses. The results are expressed as mean \pm SD in the text and as mean \pm SE in the Figures.

3 Results

3.1 Step incremental tests

A significantly greater ($p < 0.001$; Cohen's $d = 2.78$) PPO was found in Arm+leg_{INC} (337 ± 32 W) vs. Leg_{INC} (292 ± 28 W) (Figure 2A). The average relative contribution of the arms to the total power output was $22\% \pm 2\%$ in the Arm+leg_{INC} test. A higher pedaling cadence ($p < 0.01$) was found in the Leg_{INC} (84 ± 6 rpm) compared to that of the arms (76 ± 6 rpm) and legs (77 ± 7 rpm) of the Arm+leg_{INC} test. The two tests (Arm+leg_{INC} vs. Leg_{INC}) showed significant differences ($p < 0.043$) in the end-test values of $\dot{V}O_2$ ($3,913 \pm 378$ vs. $3,610 \pm 310$ mL min^{-1}), HR (186 ± 11 vs. 181 ± 10 beats min^{-1}), and V_T (2.92 ± 0.42 vs. 2.84 ± 0.40 L). When comparing the time course of the physiological and perceptual responses between Arm+leg_{INC} and Leg_{INC}, a significant ($p < 0.001$) condition \times time interaction was observed for f_R ($\eta_p^2 = 0.75$), \dot{V}_E ($\eta_p^2 = 0.74$), $\dot{V}CO_2$ ($\eta_p^2 = 0.48$), V_T ($\eta_p^2 = 0.22$) and P_{ETCO_2} ($\eta_p^2 = 0.71$). Statistically significant differences ($p < 0.001$) between Arm+leg_{INC} and Leg_{INC} were also found

when evaluating the time course of HR. Figure 3 shows where a simple main effect of condition was found. All the variables reported in Figure 3 showed a main effect of time ($p < 0.001$; $\eta_p^2 > 0.71$). No main effect of condition was found for any of the variables, but some showed $p < 0.1$, i.e. RPE ($p = 0.057$), \dot{V}_E ($p = 0.055$), $\dot{V}O_2$ ($p = 0.091$), and P_{ETCO_2} ($p = 0.091$). Due to technical problems, HR analysis was performed for 11 participants.

3.2 TTE tests at 90% of PPO_{Leg}

A significantly longer ($p < 0.001$; Cohen's $d = 2.27$) TTE was found in Arm+leg_{TTE90} (638 ± 154 s) vs. Leg_{TTE90} (307 ± 67 s) (Figure 2B). The average relative contribution of the arms to the total power output was $22\% \pm 2\%$ in the Arm+leg_{TTE90} test. A higher pedaling cadence ($p < 0.033$) was found in the Leg_{INC} (83 ± 7 rpm) compared to that of the arms (73 ± 13 rpm) and legs (76 ± 6 rpm) of the Arm+leg_{INC} test. The two tests (Arm+leg_{TTE90} vs. Leg_{TTE90}) showed significant differences ($p < 0.034$) in the end-test values of f_R (59 ± 10 vs. 53 ± 9 breaths min^{-1}), $\dot{V}O_2$ ($3,727 \pm 361$ vs. $3,581 \pm 236$ mL min^{-1}), $\dot{V}CO_2$ ($3,953 \pm 319$ vs. $4,209 \pm 302$ mL min^{-1}), HR (183 ± 9 vs. 175 ± 8 beats min^{-1}), V_T (2.68 ± 0.42 vs. 2.91 ± 0.40 L) and P_{ETCO_2} (31 ± 3 vs. 33 ± 3 mmHg). When comparing the time course of the physiological and perceptual responses between Arm+leg_{TTE90} and Leg_{TTE90}, a significant ($p < 0.017$) condition \times time interaction was observed for RPE ($\eta_p^2 = 0.32$), f_R ($\eta_p^2 = 0.52$), \dot{V}_E ($\eta_p^2 = 0.67$), $\dot{V}CO_2$ ($\eta_p^2 = 0.60$), HR ($\eta_p^2 = 0.59$) and P_{ETCO_2} ($\eta_p^2 = 0.66$). $\dot{V}O_2$ showed $p = 0.096$. Figure 4 shows where a simple main effect of condition was found. All the variables reported in Figure 4 showed a main effect of time ($p < 0.001$; $\eta_p^2 > 0.76$), while a main effect of condition ($p < 0.037$) was found for RPE ($\eta_p^2 > 0.72$), $\dot{V}CO_2$ ($\eta_p^2 > 0.40$) and V_T ($\eta_p^2 > 0.34$); $p = 0.088$ was found for HR.

3.3 TTE tests at 75% of PPO_{Leg}

A significantly longer ($p < 0.001$; Cohen's $d = 1.53$) TTE was found in Arm+leg_{TTE75} ($1,675 \pm 525$ s) vs. Leg_{TTE75} (880 ± 363 s)




FIGURE 3

Group mean response of ratings of perceived exertion (A), respiratory frequency (B), tidal volume (C), minute ventilation (D), oxygen uptake (E), carbon dioxide output (F), heart rate (G) and end-tidal partial pressure of carbon dioxide (H) for Arm+leg_{INC} (blue circles) and Leg_{INC} (grey circles). ‡*p* < 0.05 vs. Arm+leg_{INC}, **p* < 0.05 vs. Leg_{INC}.

(Figure 2C). The average relative contribution of the arms to the total power output was $21\% \pm 1\%$ in the Arm+leg_{TTE75} test. A higher pedaling cadence ($p < 0.016$) was found in the Leg_{INC} (80 ± 7 rpm) compared to that of the arms (73 ± 8 rpm) and legs (75 ± 6 rpm) of the Arm+leg_{INC} test. The two tests (Arm+leg_{TTE75} vs. Leg_{TTE75}) also showed significant differences ($p < 0.042$) in the end-test values of HR (182 ± 10 vs. 175 ± 13 beats min^{-1}) and V_T (2.40 ± 0.30 vs. 2.56 ± 0.40 L), while $p = 0.084$ was found for the end-test values of \dot{V}_E (126 ± 11 vs. 132 ± 14 L min^{-1}) and $\dot{V}CO_2$ ($3,321 \pm 270$ vs. $3,465 \pm 384$ mL min^{-1}); no significant differences ($p = 0.57$) were found for the end-test values of $\dot{V}O_2$ ($3,342 \pm 348$ and $3,379 \pm 294$ mL min^{-1}). When comparing the time course of the physiological and perceptual

responses between Arm+leg_{TTE75} and Leg_{TTE75}, a significant ($p < 0.003$) condition \times time interaction was observed for f_R ($\eta_p^2 = 0.41$), \dot{V}_E ($\eta_p^2 = 0.67$), $\dot{V}O_2$ ($\eta_p^2 = 0.35$), $\dot{V}CO_2$ ($\eta_p^2 = 0.39$), and P_{ETCO_2} ($\eta_p^2 = 0.65$). RPE showed $p = 0.068$. Statistically significant differences ($p < 0.001$) between Arm+leg_{TTE75} and Leg_{TTE75} were also found when evaluating the time course of HR. Figure 5 shows where a simple main effect of condition was found. All the variables reported in Figure 5 showed a main effect of time ($p < 0.021$; $\eta_p^2 > 0.31$), while a main effect of condition ($p < 0.011$) was found for RPE ($\eta_p^2 = 0.77$), \dot{V}_E ($\eta_p^2 = 0.48$), V_T ($\eta_p^2 = 0.53$) and P_{ETCO_2} ($\eta_p^2 = 0.47$). Due to technical problems, HR analysis was performed for 11 participants.




FIGURE 4

Group mean response of ratings of perceived exertion (A), respiratory frequency (B), tidal volume (C), minute ventilation (D), oxygen uptake (E), carbon dioxide output (F), heart rate (G) and end-tidal partial pressure of carbon dioxide (H) for Arm+leg_{TTE90} (blue circles) and Leg_{TTE90} (grey circles). †*p* < 0.05 vs. Arm+leg_{TTE90}, **p* < 0.05 vs. Leg_{TTE90}, § main effect of condition.

3.4 The performance tests considered together

When the performance tests were considered together, a significant correlation was found between f_R and RPE ($p < 0.001$; $r = 0.75$), HR and RPE ($p < 0.001$; $r = 0.69$), and \dot{V}_E and RPE ($p < 0.001$; $r = 0.80$). A graphical representation of the correlations between these variables is depicted in Figure 6.

Figure 7 shows the average response of the group when expressing \dot{V}_E as a function of V_T values, and \dot{V}_E , V_T and f_R as a function of $\dot{V}CO_2$ values. Note that the inflection point in the \dot{V}_E - V_T

relationship occurs at different V_T values, especially when comparing Arm+leg_{TTE75} and Leg_{TTE75} with the other four performance tests. A clear dissociation between f_R and $\dot{V}CO_2$ responses is observed for f_R values above 40 breaths min^{-1} .

Figure 8 shows the individual responses of \dot{V}_E , V_T and f_R expressed as a function of $\dot{V}CO_2$ values for three participants showing substantially different breathing patterns. The comparison between the responses of the three participants outlines how \dot{V}_E is more closely associated with $\dot{V}CO_2$ than V_T and f_R , and that higher values of f_R for a given $\dot{V}CO_2$ result in higher \dot{V}_E values.




FIGURE 5

Group mean response of ratings of perceived exertion (A), respiratory frequency (B), tidal volume (C), minute ventilation (D), oxygen uptake (E), carbon dioxide output (F), heart rate (G) and end-tidal partial pressure of carbon dioxide (H) for Arm+leg_{TTE75} (blue circles) and Leg_{TTE75} (grey circles). †*p* < 0.05 vs. Arm+leg_{TTE75}; **p* < 0.05 vs. Leg_{TTE75}; ‡ main effect of condition.

4 Discussion

This study aimed to systematically assess whether exercise tolerance improves with arm+leg cycling vs. leg cycling and whether f_R and perceived exertion are sensitive to the expected differences in exercise tolerance. This goal was achieved by comparing the two exercise modalities using three performance tests and two exercise paradigms (i.e. incremental test and TTE test). The main findings of the study are as follows: 1) exercise tolerance was substantially improved in all the Arm+leg tests; 2) perceived exertion, minute ventilation and respiratory frequency were particularly sensitive to the between-modality changes in exercise

tolerance observed, unlike $\dot{V}O_2$ and heart rate. These findings support the notion that respiratory frequency is a marker of physical effort during high-intensity exercise and that its time course reflects changes in exercise tolerance. This holds true even during arm+leg cycling, which is an exercise modality where the responses of $\dot{V}O_2$ and heart rate do not reflect the reduction in physical effort and the improvement in exercise tolerance observed when comparing it with leg cycling.

Our findings provide convincing evidence that arm+leg cycling substantially increases exercise tolerance when compared to (conventional) leg cycling, hence expanding on the limited literature dealing with performance differences between the two




FIGURE 6 Correlations between RPE and respiratory frequency (A), RPE and minute ventilation (B), and RPE and heart rate (C) for the Arm+leg_{INC} (blue circles), Leg_{INC} (grey circles), Arm+leg_{TTE90} (open triangles), Leg_{TTE90} (black triangles), Arm+leg_{TTE75} (yellow squares) and Leg_{TTE75} (red squares). Each symbol represents the mean value of all participants at each percentage of the TTE.





FIGURE 7 Relationship between minute ventilation and tidal volume (A), minute ventilation and carbon dioxide output (B), tidal volume and carbon dioxide output (C), and respiratory frequency and carbon dioxide output (D) for the Arm+leg_{INC} (blue circles), Leg_{INC} (grey circles), Arm+leg_{TTE90} (open triangles), Leg_{TTE90} (black triangles), Arm+leg_{TTE75} (yellow squares) and Leg_{TTE75} (red squares). Each symbol represents the mean value of all participants at each percentage of the TTE.

exercise modalities (Secher et al., 1974; Bergh et al., 1976; Nagle et al., 1984). The apparent difference in the percentage improvement in exercise tolerance found between the incremental tests and the TTE tests is, in fact, in line with the different characteristics of the two performance paradigms. Indeed, a 1% improvement in power output in an incremental test, or in a time trial, results in a performance improvement that can exceed 10% in a TTE test (Hopkins et al., 1999). Hence, the average increase in power output of about 15% that we found in the incremental test is compatible with the average increase in TTE found in the TTE

tests (i.e. 108% in the Arm+leg_{TTE90} and 90% in the Arm+leg_{TTE75}). Such an improvement in exercise tolerance observed in the Arm+leg tests implies that the effort required to sustain a given power output is substantially lower at isotime compared to that of the Leg tests, and this premise is supported by our findings.

The notion that physical effort was lower in the Arm+leg tests than in the Leg tests is substantiated by the lower values of perceived exertion and f_R generally found at isotime. While the decrease in perceived exertion during incremental Arm+leg did not reach statistical significance ($p = 0.057$), it was substantial when



comparing Arm+leg vs. Leg during the TTE tests. Indeed, perceived exertion was found to be among the most sensitive variables to variations in exercise tolerance during the TTE tests. These findings extend previous findings showing a lower RPE during arm+leg cycling vs. leg cycling at submaximal work rates (Hoffman et al., 1996; Hill et al., 2018). At isotime, f_R was significantly lower ($p < 0.003$; $\eta_p^2 > 0.40$) in the three Arm+leg tests than in the Leg tests, although this difference generally reached statistical significance in the last 20% of the Leg tests. This is an important feature of the f_R response that might have been missed in previous studies that did not compare Arm+leg and Leg until exhaustion (Robertson et al., 1986). Likewise, the reduction in f_R that is observed in the last part of a TTE test when exercise tolerance improves may not appear if the variability in TTE is not addressed on an individual basis when analyzing TTE data (Nicolò et al., 2019). We have overcome this problem by using the previously described “individual isotime”

analysis (Nicolò et al., 2019), which reduces extensively the data loss that occurs when using more traditional analyses. Our findings collectively suggest that the time course of f_R reflects changes in exercise tolerance both during incremental exercise and TTE exercise, thus supporting the study hypothesis.

The association found between f_R and RPE and the sensitivity of f_R to changes in exercise tolerance support a substantial modulation of f_R by central command (Nicolò and Sacchetti, 2023). However, a partial dissociation was found between f_R and RPE, which may suggest that also muscle afferent feedback contributed to the f_R response. Although evidence suggests that muscle afferent feedback has a greater relative contribution to f_R during moderate exercise than during high-intensity exercise (Amann et al., 2010; Dempsey et al., 2014; Girardi et al., 2021; Nicolò and Sacchetti, 2023), arm+leg cycling implies the simultaneous use of the muscle mass of both the upper and lower limbs, possibly resulting in a greater magnitude of

muscle afferent feedback, and especially of its mechanosensitive component (i.e. mechanoreflex). This may explain the slightly higher f_R shown in the first part of the TTE tests in the Arm+leg vs. the Leg modality. Indeed, it has been suggested that the relative contribution of muscle afferent feedback to ventilation is higher when the muscle mass recruited is larger (Amann et al., 2011; Dempsey et al., 2014; Nicolò and Sacchetti, 2023). Conversely, it is conceivable that the magnitude of the metabosensitive component of muscle afferent feedback (i.e. metaboreflex) was reduced at isotime in the Arm+leg tests because of the lower intramuscular metabolic perturbation. Hence, the metaboreflex cannot be ruled out as an input contributing to the decrease in f_R observed in the Arm+leg tests. Nevertheless, the relative contribution of muscle afferent feedback may reduce over time during a TTE test because the contribution of other inputs increases substantially (e.g. central command) (Dempsey et al., 2014; Nicolò and Sacchetti, 2023). We cannot exclude that afferent feedback from pulmonary mechanoreceptors or alterations in chest wall mechanics might have contributed to the partial dissociation observed between the f_R and RPE responses. Arm movements may increase the contribution to ventilation of afferent feedback from pulmonary mechanoreceptors or alter the mechanics of breathing, although these propositions require further investigation. Furthermore, it cannot be excluded that the lower pedaling cadence observed in the Arm+leg tests might have contributed to the between-modality differences observed in f_R , although variations in pedaling cadence and f_R might not be proportional, especially during high-intensity exercise (Girardi et al., 2021). Conversely, it is less plausible that metabolic acidosis or other metabolic inputs might have provided a substantial direct contribution to the f_R modulation, and the reader is referred to previous studies where evidence supporting this proposition has been reviewed (Nicolò et al., 2020a; Nicolò and Sacchetti, 2023).

The ventilatory responses observed in this study can be interpreted in the light of a recently proposed model of ventilatory control during exercise, which suggests that f_R and V_T are modulated to a large extent by behavioral and metabolic inputs respectively (Nicolò and Sacchetti, 2023). While it was proposed that ventilation is differentially regulated during incremental exercise and TTE exercise (Sybbalo et al., 1994), our findings suggest that the breathing pattern is affected by the magnitude of the inputs modulating ventilation rather than by the type of exercise paradigm. Sybbalo et al. (1994) observed a more rapid and shallow breathing pattern during a TTE test at about 76% of the PPO reached in an incremental exercise than during this latter test. The lower V_T reported by Sybbalo et al. (1994) during the TTE test is in line with the V_T response observed during the TTE tests at 75% of the PPO_{Leg} in this study. However, the breathing pattern we observed in the TTE tests at 90% of the PPO_{Leg} was, conversely, more similar to that found during the incremental exercise than during the TTE tests at 75% of the PPO_{Leg} (see Figure 7). Hence, the rapid and shallow breathing pattern is not a feature of TTE exercise, which is in contrast with what Sybbalo et al. (1994) had suggested. Conversely, the observed findings can be explained by the differential control of f_R and V_T (Nicolò et al., 2017a; Nicolò et al., 2018; Nicolò and Sacchetti, 2019; Nicolò and Sacchetti, 2023). While f_R generally shows similar peak values during incremental exercise and constant work rate exercise (Sybbalo et al., 1994; Nicolò and Sacchetti, 2023),

the V_T peak reached during exercise is largely influenced by the magnitude of metabolic inputs and is generally associated with the $\dot{V}CO_2$ peak (Nicolò and Sacchetti, 2023). As such, we found considerably lower V_T and $\dot{V}CO_2$ peak values in the Arm+leg_{TTE75} and Leg_{TTE75} tests than in the incremental tests. Conversely, when the difference in $\dot{V}CO_2$ peak between the incremental tests and the TTE tests was greatly reduced (i.e. when comparing the incremental tests with the Arm+leg_{TTE90} and Leg_{TTE90}), the difference in V_T peak values decreased accordingly. Although an association between V_T peak and $\dot{V}CO_2$ peak is commonly found when considering different exercise conditions, populations and levels of exercise capacity (Nicolò and Sacchetti, 2023), the relationship between V_T and $\dot{V}CO_2$ is not always proportional because the V_T response is to some extent influenced by the f_R response (see discussion below). Furthermore, we acknowledge that V_T is influenced by various metabolic inputs that have not been measured in this study, including metabolic acidosis (Nicolò and Sacchetti, 2023).

The association between $\dot{V}CO_2$ and V_T helps explain why \dot{V}_E resulted to be more sensitive than f_R to changes in exercise tolerance in this study. Indeed, both the magnitude of central command and that of metabolic inputs were probably higher in the Leg tests at isotime, thus increasing f_R and V_T respectively. However, the interpretation of the f_R and V_T responses observed requires careful consideration of the interdependence between the two components of \dot{V}_E , which has been advocated to explain the close match between alveolar ventilation and metabolic requirements (Haouzi, 2014). Substantial evidence suggests that V_T is fine-tuned based on f_R levels and the magnitude of metabolic inputs (Nicolò and Sacchetti, 2023), and this notion is reinforced by the present findings. While Figure 7 generally shows a consistent increase in V_T with increases in $\dot{V}CO_2$, the responses of the two variables diverge (i.e. V_T stabilizes or even decreases) when f_R starts to increase at a much steeper rate compared to $\dot{V}CO_2$. Notably, the V_T plateau did not occur at specific values of V_T or $\dot{V}CO_2$, and this is especially evident when considering the TTE tests at 75% of PPO_{Leg}, where the steeper increase in f_R occurred at relatively low $\dot{V}CO_2$ levels. Although it has been proposed that the V_T plateau that occurs during high-intensity exercise depends on mechanical constraints (Jensen et al., 1980), evidence suggesting this proposition is scarce (Nicolò et al., 2020a; Nicolò and Sacchetti, 2023). Conversely, evidence suggesting that the stabilization of V_T depends to a large extent on the increase in f_R is substantial (Nicolò et al., 2020a; Nicolò and Sacchetti, 2023), and it is even more convincing during TTE tests performed at relatively low intensities, where pulmonary mechanical limitations in healthy individuals may not occur (Nicolò et al., 2020a; Nicolò and Sacchetti, 2023).

Individual responses further suggest that V_T may not change proportionally to $\dot{V}CO_2$ values because it is affected by f_R values. At given $\dot{V}CO_2$ levels, individuals with lower f_R values show higher V_T values and *vice versa* (see Figure 8). Different combinations of f_R and V_T may guarantee the match between alveolar ventilation and metabolic requirements, and the \dot{V}_T fine-tuning feature is supposed to facilitate such a link (Nicolò and Sacchetti, 2023). Hence, individual responses reveal that \dot{V}_E is more closely associated with $\dot{V}CO_2$ than V_T , as also found in other exercise protocols (Nicolò et al., 2018; Girardi et al., 2021). The ability of the

ventilatory control system to adjust V_T according to changes in f_R has nicely been shown both at rest and during exercise in studies replacing spontaneous breathing with different levels of voluntarily imposed f_R (Lamb et al., 1965; Kennard and Martin, 1984; Haouzi and Bell, 2009; Ohashi et al., 2013; Nicolò and Sacchetti, 2023). Conversely, the ventilatory control system appears not to match metabolic requirements effectively when V_T is imposed and f_R is free to vary (Ohashi et al., 2013). In this perspective, V_T may counteract interindividual differences in f_R and guarantee an appropriate match between alveolar ventilation and $\dot{V}O_2$ for any values of f_R (Nicolò and Sacchetti, 2023). Our findings reveal the potential of comparing different exercise modalities and paradigms to shed light on the f_R and V_T modulation during high-intensity exercise.

The cardiocirculatory adjustments that occur when exercising simultaneously with the upper and lower limbs may provide further mechanistic support to the improvement in exercise tolerance observed in the Arm+leg modality. Arm+leg cycling may result in a greater peak cardiac output than Leg cycling (Secher et al., 1974; Reybrouck et al., 1975), and the higher $\dot{V}O_{2peak}$ found in the Arm+leg_{INC} vs. Leg_{INC} is in line with this notion. Hence, it is conceivable that during Arm+leg_{TTE90} and Arm+leg_{TTE75} participants were exercising at a lower fraction of peak cardiac output compared to Leg_{TTE90} and Leg_{TTE75} tests respectively, especially when similar $\dot{V}O_2$ values were found across conditions at isotime. This may have contributed to accommodating the blood flow requests of both arm and leg muscles, thus improving muscle perfusion. Indeed, the reduction in the leg power output observed in the Arm+leg TTE tests vs. the Leg TTE tests of about 20% may have reduced the leg blood flow demand in the Arm+leg modality. In turn, the lower demand of the legs may have delayed the development of leg muscle fatigue and the increase in the magnitude of central command, thus contributing to the improvement in exercise tolerance observed in the Arm+leg modality. While it has been shown that the addition of (intense) arm work to leg work reduces the leg blood flow observed at a given leg power output (Secher et al., 1977; Secher and Volianitis, 2006), the relatively low intensity sustained by the arms in our study may have not impaired leg blood flow substantially.

The fact that f_R and \dot{V}_E are considerably more sensitive to changes in exercise tolerance than $\dot{V}O_2$ and HR is particularly evident from the present study. Neither $\dot{V}O_2$ nor HR showed lower values in the Arm+leg tests than in the Leg tests at isotime, despite the lower physical effort and the improved exercise tolerance found in the Arm+leg modality. This is not surprising considering that $\dot{V}O_2$ is to a large extent associated with absolute power output during endurance cycling, although it also depends on other factors, including metabolic efficiency, which is lower for arm cycling compared to leg cycling (Cotes et al., 1969; Vokac et al., 1975; Louhevaara et al., 1990; Itoh et al., 2002). However, only a relatively small portion of the total power output is sustained by the arms during arm+leg cycling, and the oxygen uptake of arm+leg cycling has been reported to be minimally higher than that of leg cycling for a given power output (Hoffman et al., 1996). In our study, the addition of arm work to leg work resulted in a slightly higher or similar $\dot{V}O_2$ in the Arm+leg tests compared to the Leg tests. Likewise, similar or slightly higher values were observed for HR in the Arm+leg

modality at isotime, and a higher maximal HR was observed in all the Arm+leg tests. In line with our findings, previous studies had raised concerns about HR monitoring during arm+leg cycling because of the different values of maximal HR and HR relative to RPE/ $\dot{V}O_{2peak}$ that are observed when this modality is compared to leg cycling (Kitamura et al., 1981; Hoffman et al., 1996). Hence, the prescription and monitoring of arm+leg cycling should take this HR response into consideration and may benefit from the concomitant measurement of breathing variables (especially f_R), which is technically feasible even in applied settings (Massaroni et al., 2019; Nicolò et al., 2020b).

The between-modality comparison of exercise tolerance, perceived exertion and $\dot{V}O_2$ shows important practical implications of exercising in the Arm+leg modality. Our findings suggest that this exercise modality allows individuals to nearly double the amount of time spent at a given $\dot{V}O_2$ during constant work rate exercise, or, by extension, to exercise at a higher $\dot{V}O_2$ for the same exercise duration and perceived exertion. This consideration is particularly relevant for exercising individuals interested in maximizing energy expenditure, for those willing to maximize the cardiometabolic stimulus of exercise, and for those interested in lowering effort for a given absolute cardiometabolic stimulus (Hill et al., 2018). Indeed, arm+leg cycling has the potential to increase exercise adherence because a high perceived exertion is commonly viewed as one of the main barriers to exercise participation (Cheval and Boissongier, 2021), and a relatively low perceived exertion may be associated with a sufficient cardiometabolic stimulus in this exercise modality. Furthermore, considering that exercise tolerance is closely associated with morbidity and mortality (Kokkinos et al., 2010; Nesti et al., 2020), arm+leg cycling may have clinical implications. Arm+leg cycling involves the simultaneous use of arm and leg muscles and may result in a time-efficient training strategy for enhancing both health and performance (Zinner et al., 2017). However, further research is needed to test these propositions.

It is worth mentioning that the Arm+leg modality poses some measurement challenges that we had to face in this study. First, we did not attempt to compute the GET and the RCP because of the relatively high initial stage of the incremental tests imposed by the ergometer (i.e. 150 W), which has limited their detection. As such, the TTE tests were not prescribed based on exercise intensity domains. Hence, especially in the Arm+leg_{TTE75} test, it is possible that some participants were exercising in the severe intensity domain while others were in the heavy domain. Second, the measurement of blood lactate is very challenging during the Arm+leg modality, and we decided not to collect it to avoid interfering with the performance tests. Third, perceived exertion can only be rated verbally during the Arm+leg tests, thus generating breathing artifacts and affecting gas exchange measures. However, we used a filtering technique (Lamarra et al., 1987) that addressed this limitation and helped preserve the integrity of the physiological responses. Even the other limitations outlined were partially counteracted by the experimental design and the method of analysis used. Indeed, we performed a detailed between-modality comparison of the responses of some of the main physiological variables commonly used to compute the GET and the RCP. This comparison reveals a

greater metabolic perturbation at isotime in the Leg tests, especially when considering the time courses of \dot{V}_E , $\dot{V}CO_2$ and P_{ETCO_2} . Hence, it is conceivable that participants were exercising at a lower relative exercise intensity during the Arm+leg tests, not only from an effort perspective but also from a metabolic perspective.

5 Conclusion

This study shows that exercise tolerance is substantially higher in the Arm+leg modality than in the leg cycling modality. The average improvement in exercise tolerance was 15% in the incremental test and 108% and 90% in the TTE tests at 90% and 75% of PPO_{Leg} respectively. Perceived exertion, minute ventilation and respiratory frequency were among the most sensitive variables to the improvement in exercise tolerance provided by the Arm+leg modality, hence suggesting that the common mechanism modulating these variables (i.e. central command) plays an important role in endurance performance. These findings reinforce the notion that respiratory frequency is a better marker of physical effort than oxygen uptake and heart rate. Our results have implications for devising exercise strategies to reduce perceived exertion while maintaining a high cardiometabolic demand, and for maximizing energy expenditure for the same level of effort exerted.

Data availability statement

The raw data supporting the conclusion of this article will be made available by the authors, without undue reservation.

Ethics statement

The studies involving human participants were reviewed and approved by Institutional Review Board of the University of Rome “Foro Italico.” The patients/participants provided their written informed consent to participate in this study.

References

- Amann, M., Blain, G. M., Proctor, L. T., Sebranek, J. J., Pegelow, D. F., and Dempsey, J. A. (2010). Group III and IV muscle afferents contribute to ventilatory and cardiovascular response to rhythmic exercise in humans. *J. Appl. Physiol.* 109, 966–976. doi:10.1152/jappphysiol.00462.2010
- Amann, M., Runnels, S., Morgan, D. E., Trinity, J. D., Fjeldstad, A. S., Wray, D. W., et al. (2011). On the contribution of group III and IV muscle afferents to the circulatory response to rhythmic exercise in humans. *J. Physiology* 589, 3855–3866. doi:10.1113/jphysiol.2011.209353
- Astrand, P. O., and Saltin, B. (1961). Maximal oxygen uptake and heart rate in various types of muscular activity. *J. Appl. Physiol.* 16, 977–981. doi:10.1152/jappl.1961.16.6.977
- Bergevin, M., Steele, J., Payen de la Garanderie, M., Feral-Basin, C., Marcora, S. M., Rainville, P., et al. (2023). Pharmacological blockade of muscle afferents and perception of effort: A systematic review with meta-analysis. *Sports Med.* 53, 415–435. doi:10.1007/s40279-022-01762-4
- Bergh, U., Kanstrup, I. L., and Ekblom, B. (1976). Maximal oxygen uptake during exercise with various combinations of arm and leg work. *J. Appl. Physiol.* 41, 191–196. doi:10.1152/jappl.1976.41.2.191
- Bland, J. M., and Altman, D. G. (1995). Calculating correlation coefficients with repeated observations: Part 1-Correlation within subjects. *BMJ* 310, 446. doi:10.1136/bmj.310.6977.446
- Borg, G. (1998). Borg’s perceived exertion and pain scales. *Hum. Kinet.* 30, 1461. doi:10.1249/00005768-199809000-00018
- Cheval, B., and Boisgontier, M. P. (2021). The theory of effort minimization in physical activity. *Exerc Sport Sci. Rev.* 49, 168–178. doi:10.1249/JES.0000000000000252
- Cohen, J. (1988). *Statistical power analysis for the behavioural sciences* 2nd ed.
- Cotes, J. E., Allsopp, D., and Sardi, F. (1969). Human cardiopulmonary responses to exercise: Comparisons between progressive and steady state exercise, between arm and leg exercise, and between subjects differing in body weight. *Q. J. Exp. Physiol. Cogn. Med. Sci.* 54, 211–222. doi:10.1113/expphysiol.1969.sp002019
- Dempsey, J. A., Blain, G. M., and Amann, M. (2014). Are type III-IV muscle afferents required for a normal steady-state exercise hyperpnoea in humans? *J. Physiology* 592, 463–474. doi:10.1113/jphysiol.2013.261925
- Girardi, M., Nicolò, A., Bazzucchi, I., Felici, F., and Sacchetti, M. (2021). The effect of pedalling cadence on respiratory frequency: Passive vs. active exercise of different intensities. *Eur. J. Appl. Physiol.* 121, 583–596. doi:10.1007/s00421-020-04533-z
- Gleser, M. A., Hoestman, D. H., and Mello, R. P. (1974). The effect on $\dot{V}O_2$ max of adding arm work to maximal leg work. *Med. Sci. Sports Exerc* 6, 104–107. doi:10.1249/00005768-197400620-00018

Author contributions

AN, MS, and FF contributed to the conception and design of the study. MG performed data collection. MG and AN analyzed data. AN wrote the first draft of the manuscript. AN, MG, IB, MS, and FF contributed to data interpretation. All authors contributed to the article and approved the submitted version.

Funding

The study was funded by Harken Italy SPA (Limido Comasco, Como). AN was supported by a Post-Doc Research Grant funded by Harken Italy SPA (Limido Comasco, Como). The article publication fees were provided by the University of Rome “Foro Italico” (CDR2. BANDO2022NA).

Acknowledgments

The authors would like to thank Andrea Merello and Michele Cazzaro from Harken Italy SPA for their enthusiasm and valuable help in the different stages of the project.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Publisher’s note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

- Haouzi, P., and Bell, H. J. (2009). Control of breathing and volitional respiratory rhythm in humans. *J. Appl. Physiol.* 106, 904–910. doi:10.1152/jappphysiol.90675.2008
- Haouzi, P. (2014). Tracking pulmonary gas exchange by breathing control during exercise: Role of muscle blood flow. *J. Physiology* 592, 453–461. doi:10.1113/jphysiol.2013.261396
- Hill, M., Talbot, C., Puddiford, M., and Price, M. (2018). Cardiorespiratory and perceptual responses to self-regulated and imposed submaximal arm–leg ergometry. *Eur. J. Appl. Physiol.* 118, 1011–1019. doi:10.1007/s00421-018-3838-7
- Hoffman, M. D., Kassay, K. M., Zeni, A. I., and Clifford, P. S. (1996). Does the amount of exercising muscle alter the aerobic demand of dynamic exercise? *Eur. J. Appl. Physiol. Occup. Physiol.* 74, 541–547. doi:10.1007/BF02376770
- Hopkins, W. G., Hawley, J. A., and Burke, L. M. (1999). Design and analysis of research on sport performance enhancement. *Med. Sci. Sports Exerc* 31, 472–485. doi:10.1097/00005768-199903000-00018
- Itoh, M., Fukuoka, Y., Grassi, B., Marconi, C., Cerretelli, P., Araki, H., et al. (2002). VE response to VCO₂ during exercise is unaffected by exercise training and different exercise limbs. *Jpn. J. Physiology* 52, 489–496. doi:10.2170/jjphysiol.52.489
- Jensen, J. L., Lyager, S., and Pedersen, O. F. (1980). The relationship between maximal ventilation, breathing pattern and mechanical limitation of ventilation. *J. Physiol.* 309, 521–532. doi:10.1113/jphysiol.1980.sp013524
- Kennard, C. D., and Martin, B. J. (1984). Respiratory frequency and the oxygen cost of exercise. *Eur. J. Appl. Physiol. Occup. Physiol.* 52, 320–323. doi:10.1007/BF01015218
- Kitamura, K., Yamaji, K., and Shephard, R. J. (1981). Heart rate predictions of exercise intensity during arm, leg and combined arm/leg exercise. *J. Hum. Ergol. (Tokyo)* 10, 151–160.
- Kokkinos, P., Myers, J., Faselis, C., Panagiotakos, D. B., Doumas, M., Pittaras, A., et al. (2010). Exercise capacity and mortality in older men: A 20-year follow-up study. *Circulation* 122, 790–797. doi:10.1161/CIRCULATIONAHA.110.938852
- Lamarra, N., Whipp, B. J., Ward, S. A., and Wasserman, K. (1987). Effect of interbreath fluctuations on characterizing exercise gas exchange kinetics. *J. Appl. Physiol.* 62, 2003–2012. doi:10.1152/jappphysiol.1987.62.5.2003
- Lamb, T. W., Anthonisen, N. R., and Tenney, S. M. (1965). Controlled frequency breathing during muscular exercise. *J. Appl. Physiol.* 20, 244–248. doi:10.1152/jappphysiol.1965.20.2.244
- Louhevaara, V., Sovijarvi, A., Ilmarinen, J., and Teraslinna, P. (1990). Differences in cardiorespiratory responses during and after arm crank and cycle exercise. *Acta Physiol. Scand.* 138, 133–143. doi:10.1111/j.1748-1716.1990.tb08825.x
- Marcora, S. (2009). Perception of effort during exercise is independent of afferent feedback from skeletal muscles, heart, and lungs. *J. Appl. Physiol.* 106, 2060–2062. doi:10.1152/jappphysiol.90378.2008
- Massaroni, C., Nicolò, A., Lo Presti, D., Sacchetti, M., Silvestri, S., and Schena, E. (2019). Contact-based methods for measuring respiratory rate. *Sensors* 19, 908. doi:10.3390/s19040908
- Nagle, F. J., Richie, J. P., and Giese, M. D. (1984). $\dot{V}O_2$ max responses in separate and combined arm and leg air-braked ergometer exercise. *Med. Sci. Sports Exerc* 16, 563–566. doi:10.1249/00005768-198412000-00007
- Nesti, L., Pugliese, N. R., Sciuto, P., and Natali, A. (2020). Type 2 diabetes and reduced exercise tolerance: A review of the literature through an integrated physiology approach. *Cardiovasc Diabetol.* 19, 134. doi:10.1186/s12933-020-01109-1
- Nicolò, A., Bazzucchi, I., Haxhi, J., Felici, F., and Sacchetti, M. (2014). Comparing continuous and intermittent exercise: An “isoeffort” and “isotime” approach. *PLoS One* 9, e94990. doi:10.1371/journal.pone.0094990
- Nicolò, A., Girardi, M., Bazzucchi, I., Felici, F., and Sacchetti, M. (2018). Respiratory frequency and tidal volume during exercise: Differential control and unbalanced interdependence. *Physiol. Rep.* 6, e13908. doi:10.14814/phy2.13908
- Nicolò, A., Marcora, S. M., Bazzucchi, I., and Sacchetti, M. (2017a). Differential control of respiratory frequency and tidal volume during high-intensity interval training. *Exp. Physiol.* 102, 934–949. doi:10.1113/EP086352
- Nicolò, A., Marcora, S. M., and Sacchetti, M. (2016). Respiratory frequency is strongly associated with perceived exertion during time trials of different duration. *J. Sports Sci.* 34, 1199–1206. doi:10.1080/02640414.2015.1102315
- Nicolò, A., Marcora, S. M., and Sacchetti, M. (2020a). Time to reconsider how ventilation is regulated above the respiratory compensation point during incremental exercise. *J. Appl. Physiol.* 128, 1447–1449. doi:10.1152/jappphysiol.00814.2019
- Nicolò, A., Massaroni, C., and Passfield, L. (2017b). Respiratory frequency during exercise: The neglected physiological measure. *Front. Physiol.* 8, 922. doi:10.3389/fphys.2017.00922
- Nicolò, A., Massaroni, C., Schena, E., and Sacchetti, M. (2020b). The importance of respiratory rate monitoring: From healthcare to sport and exercise. *Sensors Switz.* 20, 6396. doi:10.3390/s20216396
- Nicolò, A., and Sacchetti, M. (2019). A new model of ventilatory control during exercise. *Exp. Physiol.* 104, 1331–1332. doi:10.1113/EP087937
- Nicolò, A., and Sacchetti, M. (2023). Differential control of respiratory frequency and tidal volume during exercise. *Eur. J. Appl. Physiol.* 123, 215–242. doi:10.1007/s00421-022-05077-0
- Nicolò, A., Sacchetti, M., Girardi, M., McCormick, A., Angius, L., Bazzucchi, I., et al. (2019). A comparison of different methods to analyse data collected during time-to-exhaustion tests. *Sport Sci. Health* 15, 667–679. doi:10.1007/s11332-019-00585-7
- Ohashi, S., Izumizaki, M., Atsumi, T., and Homma, I. (2013). CO₂ homeostasis is maintained in conscious humans by regulation of tidal volume, but not of respiratory rhythm. *Respir. Physiol. Neurobiol.* 186, 155–163. doi:10.1016/j.resp.2013.01.008
- Reybrouck, T., Heigenhauser, G. F., and Faulkner, J. A. (1975). Limitations to maximum oxygen uptake in arms, leg, and combined arm–leg ergometry. *J. Appl. Physiol.* 38, 774–779. doi:10.1152/jappphysiol.1975.38.5.774
- Robertson, R. J., Falkel, J. E., Drash, A. L., Swank, A. M., Metz, K. F., Spungen, S. A., et al. (1986). Effect of blood pH on peripheral and central signals of perceived exertion. *Med. Sci. Sports Exerc* 18, 114–122. doi:10.1249/00005768-198602000-00019
- Secher, N. H., Clausen, J. P., Klausen, K., Noer, I., and Trap-Jensen, J. (1977). Central and regional circulatory effects of adding arm exercise to leg exercise. *Acta Physiol. Scand.* 100, 288–297. doi:10.1111/j.1748-1716.1977.tb05952.x
- Secher, N. H., Ruberg Larsen, N., Binkhorst, R. A., and Bonde Petersen, F. (1974). Maximal oxygen uptake during arm cranking and combined arm plus leg exercise. *J. Appl. Physiol.* 36, 515–518. doi:10.1152/jappphysiol.1974.36.5.515
- Secher, N. H., and Volianitis, S. (2006). “Are the arms and legs in competition for cardiac output?,” in *Medicine and science in sports and exercise*. doi:10.1249/01.mss.0000230343.64000.ac
- Syabbalo, N. C., Krishnan, B., Zintel, T., and Gallagher, C. G. (1994). Differential ventilatory control during constant work rate and incremental exercise. *Respir. Physiol.* 97, 175–187. doi:10.1016/0034-5687(94)90024-8
- Van De Walle, G. P., and Vukovich, M. D. (2018). The effect of nitrate supplementation on exercise tolerance and performance: A systematic review and meta-analysis. *J. Strength Cond. Res.* 32, 1796–1808. doi:10.1519/jsc.0000000000002046
- Vokac, Z., Bell, H., Bautz Holter, E., and Rodahl, K. (1975). Oxygen uptake/heart rate relationship in leg and arm exercise, sitting and standing. *J. Appl. Physiol.* 39, 54–59. doi:10.1152/jappphysiol.1975.39.1.54
- Zimmer, C., Sperlich, B., Born, D. P., and Michels, G. (2017). Effects of combined high intensity arm and leg training on performance and cardio-respiratory measures. *J. Sports Med. Phys. Fit.* 57, 969–975. doi:10.23736/S0022-4707.16.06539-7